

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : A61K 39/395, 39/385, 38/00, G01N 33/53	A1	(11) International Publication Number: WO 00/20032 (43) International Publication Date: 13 April 2000 (13.04.00)
(21) International Application Number: PCT/US99/23251 (22) International Filing Date: 5 October 1999 (05.10.99) (30) Priority Data: 60/103,284 6 October 1998 (06.10.98) US (71) Applicants: TRUSTEES OF DARTMOUTH COLLEGE [US/US]; Technology Transfer Office, 11 Rope Ferry Road, Hanover, NH 03755-1404 (US). MEDAREX, INC. [US/US]; 1545 Route 22 East, Annandale, NJ 08801 (US). (72) Inventors: GUYRE, Paul, M.; 21 Pinneo Hill Road, Hanover, NH 03755 (US). GOLDSTEIN, Joel, J.; 14 Lewis Place, Piscataway, NJ 08854 (US). WU, Zining; 55 Meadow Lane, Norristown, PA 19403 (US). SUN, Wanwen; 55 Meadow Lane, Norristown, PA 19403 (US). (74) Agents: LICATA, Jane, Massey et al.; Law Offices of Jane Massey Licata, 66 E. Main Street, Marlton, NJ 08053 (US).		(81) Designated States: CA, JP, European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE). Published <i>With international search report.</i>
(54) Title: RECOMBINANT CAT ALLERGEN, Fel dI, EXPRESSED IN BACULOVIRUS FOR DIAGNOSIS AND TREATMENT OF CAT ALLERGY (57) Abstract Recombinant Fel dI cat allergens expressed in baculovirus for diagnosis and treatment of allergy to cats in humans are provided.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

- 1 -

**RECOMBINANT CAT ALLERGEN, Fel dI, EXPRESSED IN BACULOVIRUS
FOR DIAGNOSIS AND TREATMENT OF CAT ALLERGY**

Background of the Invention

Fel dI is the major allergen from cats. Natural Fel dI
5 consists of two polypeptide chains, chain 1(ch1) and chain
2(ch2) which are normally linked by a disulfide bond. Fel dI
has been cloned and sequenced. However, the immunoreactivity
of rFel dI chains expressed in bacteria is not comparable to
that of the natural allergen (Shint et al. *JACI* 1995,1221).

10 **Summary of the Invention**

An object of the present invention is to provide a
composition for diagnosis and treatment of cat allergy in
humans comprising a baculovirus expressed recombinant Fel dI.

Brief Description of the Figure

15 Figure 1 shows a schematic of the final construct of
H22-Fel dI Ch1+Ch2 in pAcSAG-LIC.

Detailed Description of the Invention

It has now been found that the immunoreactivity of rFel
dI for IgG and IgE antibody is improved dramatically by
20 expressing the allergen in baculovirus.

Recombinant Fel dI, rFel dI Ch1+Ch2, in which the two
chains are expressed in series and linked together by a
glycine/serine linker (referred to herein as H22-), and CD64-
targeted Fel dI (sFv22;Fel dI), which consists of the
25 foregoing rFel d I Ch1+Ch2 linked to the sFv of monoclonal

- 2 -

antibody H22 (mAb H22) (referred to herein as H22+) were genetically constructed. Mab H22 is the humanized anti-CD64 antibody (Graziano et al. *J Immunol.* **1995** 155, 4996-5002). Since CD64 is only expressed by monocytes and dendritic cells, it is believed that the H22+ fusion protein targets Fel dI specifically to monocytes and dendritic cells via the sFv component, which is derived from the anti-CD64 monoclonal antibody H22. The molecular weight of the H22+ and H22- were 49 kd and 22 kd, respectively.

H22+ and H22- baculovirus expressed rFel dIs were purified by Ni affinity chromatography and compared with natural Fel dI (nFel dI) by ELISA using a panel of anti-Fel dI monoclonal antibodies and by RIA binding of the antigen to human IgE and IgG antibodies. Both H22+ and H22- rFel dI proteins demonstrated similar binding to nFel dI in ELISA using different combinations of monoclonal antibodies. Results from an ELISA are depicted in the following Table 1.

Table 1:

Capture Ab	nFel dI	H22+FeldI Ch1+Ch2	rFeldI Ch1	H22+FeldI Ch1
1G9 (EPI-B, CH1)	++++	++++	++	+
8F3 (EBI-B, CH1)	+	+	-	-
2H4 (EPI-C, CH2)	+++	+++	-	-
10G7 (EPI-D, ?)	+	+	-	-
11F5 (R&A, CH1)	-	-	-	-
8H6 (R&A?, ?)	-	-	-	-
6F9 (?, CH1)	++++	++++	++++	++

- 3 -

The detection antibody in these studies was 3E4-biotin.

By inhibition RIA, H22+ rFel dI showed identical inhibition curves to nFel dI using IgG antibody in pooled sera from either Japanese (n=10) or US (n=6) cat allergic patients. The H22+ rFel dI inhibited binding of nFel dI by >95%. Excellent correlations were obtained by linear regression analysis comprising IgE antibody to H22+ rFel dI (n=155, r=0.72, p<0.001) or IgE antibody to H22- rFel dI (n=258, r=0.72, p<0.001) with nFel dI. These data show that IgG and IgE antibody binding by baculovirus expressed rFel dI is identical to nFel dI.

Accordingly, the baculovirus expressed rFel dIs of the present invention are believed to be useful in the diagnosis and treatment of cat allergy. Use of the rFel dI allergens of the present invention to diagnose a cat allergy in human serum samples is performed routinely in accordance with well known procedures. Similarly, incorporation of the allergens of the present invention into a treatment regime such as allergy shots for the treatment of cat allergies in humans is also performed in accordance with well known techniques.

The H22+ construct of the present invention is also useful in targeting of Fel dI to monocytes and dendritic cells for studies of antigen presentation and T cell responses in cat allergic patients.

The following nonlimiting examples are provided to further illustrate the present invention.

EXAMPLES

Example 1: Plasmids and oligonucleotides

Baculovirus expression vector pAcSAG-LIC was purchased from Pharmingen. H22 sFv (encoding $V_H V_L$ of the anti-CD64

- 4 -

antibody H22) was cloned from vector pJG225 (Medarex, Inc. Annandale, NJ, USA) into the *Bam*HI and *Xba*I sites of pAcSAG-LIC and renamed pTJ225. Vectors pET11dΔHR chain-1 *Feld*I and pET11dΔHR chain-2 *Feld*I were provided by Immunologic (Waltham, MA). Chain 1 of *Feld*I was cloned into pTJ225 by PCR cloning. Chain 2 was cloned into vector pCRTM2.1 of the TA cloning kit (Invitrogen, Carlsbad, CA, USA). Primers were ordered from Integrated DNA Technologies (IDT, Coralville, IA) and contained the following sequences:

10 Chain 1:

forward primer: 32 mer (SEQ ID NO:1)

5' AGG ACT CGA GTG **AAA TTT GCC CAG CCG TGA AG** 3'
 *Xho*I

backward primer: 36 mer (SEQ ID NO:2)

15 5' TAA ACT TCG CGG CCG C CA TAT GAC **ACA GAG GAC TTG** 3'
 *Not*I *Nde*I

Chain 2:

forward primer: 28 mer (SEQ ID NO:3)

20 5' GGG GCT GCA GGT **CAA GAT GGC GGA AAC T** 3'
 *Pst*I

backward primer: 33 mer (SEQ ID NO:4)

5' GTT GTC AGC AGC GGC CGC TCT **CCC CAA AGT GTT** 3'
 *Not*I

Sequences complementary to the cDNA are shown in bold.

25 To clone chain 1 and chain 2 succeedingly after H22, a linker oligo was designed. This linker oligo encodes the flexible peptide linker (Gly₄Ser)₃. Unique restriction sites were designed on both sides of the linker creating sticky ends

- 5 -

immediately after annealing. The DNA sequence of the linker is described below.

Linker:

sense, 54 mer (SEQ ID NO:5)

5 5' TATG(GGT GGA GGA GGT TCT)_{x3}CTGCA 3'
NdeI PstI

antisense, 48 mer

5' G(AGAACCTCCTCCACC)_{x3}CA 3' (SEQ ID NO:6)

To generate H22-FeldI Ch1+Ch2 in baculovirus expression vector
10 pAcSAG-LIC, FeldI Ch1 digested with *XhoI* and *NdeI*, linker with
sticky ends *NdeI* and *PstI* and FeldI Ch2 restricted with *PstI*
and *NotI* were ligated into the *XhoI* and *NotI* sites of pTJ225
in a four part ligation subcloning. The final construct is
depicted in Figure 1.

15 **Example 2: Generation of Recombinant Virus containing the
H22-FeldI Ch1+Ch2 sequences**

To generate recombinant virus, 3 x 10⁹ Sf9 cells in 60
mm tissue culture dish were co-transfected with 1 µg of
baculovirus expression plasmid containing the genes of
20 interest, using the transfection protocol according to the
manufacturer's instructions. Four days after the
transfection, the culture supernatant containing the
recombinant viruses was collected. The titers of recombinant
virus were then amplified to 5-10 x 10⁸ plaque forming units
25 (pfu)/ml by infecting more Sf9 cells.

Example 3: Protein Expression and Purification

High Five™ insect cells were chosen for large-scale
production of recombinant protein. To determine the time

- 6 -

course of recombinant protein expression , a monolayer of High Five™ cells in a T-75 culture flask was infected with high titer recombinant virus at a multiplicity of infection (MOI) of 10. At specific intervals following infection, culture supernatant was collected and the proteins were precipitated with 72% trichloroacetic acid and 0.15% sodium deoxycholate. After resuspension in 0.1 volumes of sample buffer, SDS-PAGE (10-20% gradient gel) was performed and the gel was stained with Coomassie Blue R-250. Large scale expression was accomplished by infecting large volumes of suspension cultured cells. Cell-free supernatants were harvested 72 hours post-infection by removing the cells at 1000 rpm for 10 minutes at 4°C. At this time point expression of antibody fusion protein reached its peak in cell culture supernatants while there was limited intracellular protein resulting from cell lysis. The cell-free culture supernatants were then concentrated 10-fold, dialyzed and loaded onto a nickel (Ni)-affinity column (Novagen, Inc.). After washing the loading buffer, proteins were eluted with a linear gradient of imidazole in the same buffer. Fractions containing recombinant antibody-fusion protein were pooled and dialyzed. The pooled fractions were then applied to an anion-exchange column (Econo-Pac S-cartridge, Bio-Rad). the flow-through, containing recombinant protein, was collected and dialyzed in phosphate-buffered saline (PBS). The purity of all protein preparations was monitored by SDS-PAGE and was at least 95% homogenous. Protein concentrations were determined from A280nm values calculated with molar extinction coefficient of 60293.0 A280 nm/mole. Yield was approximately 4-6 mg of purified recombinant protein per liter of Hi-5 culture supernatant.

- 7 -

What is claimed is:

1. A composition comprising a baculovirus expressed recombinant Fel dI.
2. The composition of claim 1 wherein the baculovirus
5 expressed recombinant Fel dI comprises chain 1 and chain 2
expressed in series and linked together by a glycine/serine
linker.
3. The composition of claim 2 further comprising a sFv
of monoclonal antibody H22.
- 10 4. A method of diagnosing a human with cat allergy
comprising contacting a serum sample from a human with a
composition of claim 1 and determining the immunoreactive
response of the serum sample to the composition of claim 1
wherein an immune reaction against the composition is
15 indicative of an allergy to cats.
5. A method of protecting a human against a cat
allergy comprising administering to a human a composition of
claim 1.




Fig. 1

SEQUENCE LISTING

<110> Guyre, Paul M.
Goldstein, Joel
Wu, Zining
Sun, Wanwen
Trustees of Dartmouth College
Medarex, Inc.

<120> Recombinant Cat Allergen, Fel dI, Expressed in
Baculovirus for Diagnosis and Treatment of Cat Allergy

<130> DC-0119

<140>
<141>

<150> 60/103,284
<151> 1998-10-06

<160> 6

<170> PatentIn Ver. 2.0

<210> 1
<211> 32
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:Synthetic

<400> 1
aggactcgag tgaaatttgc ccagccgtga ag 32

<210> 2
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:Synthetic

<400> 2

<210> 3
<211> 28
<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:Synthetic

<400> 3

<210> 4

<211> 33

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:Synthetic

<400> 4

<210> 5

<211> 54

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:Synthetic

<400> 5

<210> 6

<211> 48

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:Synthetic

<400> 6

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US99/23251

A. CLASSIFICATION OF SUBJECT MATTER

IPC(6) :A61K 39/395, 39/385, 38/00; G01N 33/53

US CL :424/134.1, 135.1, 178.1, 179.1, 193.1; 435/7.1; 514/2

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 424/134.1, 135.1, 178.1, 179.1, 193.1; 435/7.1; 514/2

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

Please See Extra Sheet.

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	GUYRE et al. Increased potency of Fc-receptor-targeted antigens. Cancer Immunol Immunother. 1997, Vol. 45, pages 146-148, especially Abstract, page 148, column 2 last sentence and page 148 column 1, first full paragraph.	3
Y	ROGERS et al. POTENTIAL THERAPEUTIC RECOMBINANT PROTEINS COMPRISED OF PETPIDES CONTAINING RECOMBINED T CELL EPITOPES. Molecular Immunology. 1994, Vol. 31, No. 13, pages 955-966, especially Abstract.	1-5
Y	US 5,359,045 A (SOUBRIER ET AL) 25 October 1994 (25/10/94), see entire document, especially column 7, lines 28-33.	1-5



Further documents are listed in the continuation of Box C.



See patent family annex.

* Special categories of cited documents:	*T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
A document defining the general state of the art which is not considered to be of particular relevance	*X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
E earlier document published on or after the international filing date	*Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
L document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	*Z* document member of the same patent family
O document referring to an oral disclosure, use, exhibition or other means	
P document published prior to the international filing date but later than the priority date claimed	

Date of the actual completion of the international search

20 DECEMBER 1999

Date of mailing of the international search report

21 JAN 2000

Name and mailing address of the ISA/US
Commissioner of Patents and Trademarks
Box PCT
Washington, D.C. 20231

Facsimile No. (703) 305-3230

Authorized officer

MARIANNE DIBRINO

Telephone No. (703) 308-0196

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US99/23251

C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	US 5,356,622 A (HEATH ET AL) 18 October 1994 (18/10/94), see entire document, especially column 6, lines 37-38.	1-5
Y	US 5,795,862 A (FRANK ET AL) 18 August 1998 (18/08/98), see entire document, especially column 38, lines 15-67, column 39, lines 1-26 and column 40, lines 36-51.	4, 5
Y	SANA et al. Expression and Ligand Binding Characterization of the b-Subunit (p75) Ectodomain of the Interleukin-2 Receptor. Biochemistry. 1994, Vol. 33, pages 5838-5845, especially page 5839, column 1, last sentence.	1-5
Y	RHODE et al. Single-Chain MHC Class II Molecules Induce T Cell Activation and Apoptosis. The Journal of Immunology. 1996, Vol. 157, pages 4885-4889, especially page 4889, lines 7-13.	1-5

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US99/23251

B. FIELDS SEARCHED

Electronic data bases consulted (Name of data base and where practicable terms used):

STN (BIOSCIENCE, CAPLUS, USPATFUL)

search terms: Goldstein, Joel, Wu, Zining, Sun, Wanwen, Fel dI, baculovirus, monoclonal, anti-cd64, recombinant Fel dI, rFel dI, cat allergy, chain 1, chain 2, Guyre, Paul, antigens, autoimmune, vaccination, allergens, epitopes, H22